

Mercury Pollution in Aquatic Ecosystems






Karen A. Merritt, PhD MPH



What do we know regarding the biomagnification process?

- Sulfate (SO_4^{2-}) + low-to-no dissolved oxygen (O_2) increases activity of sulfate-reducing **bacteria** (SRB);
- SRB in the presence of inorganic mercury (Hg^{2+}) generate methyl mercury (CH_3Hg^+) as an accident/**by-product of respiration**;
- Methyl mercury is **retained in biological tissue** more significantly than inorganic mercury because of the additional $-\text{CH}_3$ (methyl) group;
- **Bioaccumulation** of methyl mercury occurs because the depuration (loss) rate of methyl mercury from biological tissue is much lower/slower than the loss rate of inorganic mercury;
- **Biomagnification** happens through the **trophic transfer** of bioaccumulated methyl mercury from small prey species to larger prey (or predator) species to largest predator species (including humans)





A Global Model for Methylmercury Formation and Uptake at the Base of Marine Food Webs

Yanxu Zhang^{1,2}, Anne L. Soerensen^{3,4}, Amina T. Scharup^{5,6}, and Elsie M. Sunderland^{2,3}

¹Joint International Research Laboratory of Atmospheric and Earth System Sciences, School of Atmospheric Sciences, Nanjing University, Nanjing, China, ²Harvard John A. Paulson School of Engineering & Applied Sciences, Harvard University, Cambridge, MA, USA, ³Department of Environmental Health, Harvard T.H. Chan School of Public Health, Harvard University, Boston, MA, USA, ⁴Department of Environmental Science and Analytical Chemistry, Stockholm University, Stockholm, Sweden, ⁵ Scripps Institution of Oceanography, La Jolla, CA, USA

Abstract Monomethylmercury (CH₃Hg) is the only form of mercury (Hg) known to biomagnify in food webs. Here we investigate factors driving methylated mercury [MeHg = CH₃Hg + (CH₃)₂Hg] production and degradation across the global ocean and uptake and trophic transfer at the base of marine food webs. We develop a new global 3-D simulation of MeHg in seawater and phyto/zooplankton within the Massachusetts Institute of Technology general circulation model. We find that high modeled MeHg concentrations in polar regions are driven by reduced demethylation due to lower solar radiation and colder temperatures. In the eastern tropical subsurface waters of the Atlantic and Pacific Oceans, the model results suggest that high MeHg concentrations are associated with enhanced microbial activity and atmospheric inputs of inorganic Hg. Global budget analysis indicates that upward advection/diffusion from subsurface ocean provides 17% of MeHg in the surface ocean. Modeled open ocean phytoplankton concentrations are relatively uniform because lowest modeled seawater MeHg concentrations occur in oligotrophic regions with the smallest size classes of phytoplankton, with relatively high uptake of MeHg and vice versa. Diatoms and synchococcus are the two most important phytoplankton categories for transferring MeHg from seawater to herbivorous zooplankton, contributing 35% and 25%, respectively. Modeled ratios of MeHg concentrations between herbivorous zooplankton and phytoplankton are 0.74–0.78 for picoplankton (i.e., no biomagnification) and 2.6–4.5 for eukaryotic phytoplankton. The spatial distribution of the trophic magnification factor is largely determined by the zooplankton concentrations. Changing ocean biogeochemistry resulting from climate change is expected to have a significant impact on marine MeHg formation and bioaccumulation.



Think about **risk profiles** for different locations in terms of:

- **Physical factors** – what is the burial rate of contamination based on the geological background (i.e., how much sediment is available for burying contamination quickly to a depth below the biological mixed depth or biologically active zone?); how do hydrodynamics impact stable burial?
- **Chemical factors** – what factors are present that can create the conditions in which SRB are active? Factors of concern are those that contribute sulfate (SO_4^{2-}) and biochemical oxygen demand (BOD) such that significant O_2 consumption occurs. Factors can be anthropogenic but aren't always.
- **Biological factors** – what are the species of concern and what do trophic transfer pathways look like? For human health concerns, what are the frequency and frameworks for consumption (i.e., recreational and infrequent vs frequent and culturally or socio-economically significant)?

PHYSICAL

Sufficient sedimentation and low erosion potential to allow for stable burial



BIOLOGICAL



Multi-trophic level food chains with a top predator species that is frequently consumed



- ★ = Wabigoon River, ON
- ★ = Penobscot River, ME
- ★ =

CHEMICAL

Factors that contribute BOD and sulfate (SO_4^{2-}) and result in O_2 consumption



Think of this overview of risk profiles as describing orbits around a worst-case ecological and human health scenario

Site Comparison

Penobscot River Estuary, Maine

- Glaciated terrain and low sed. rates (~ 0.5 cm/yr)
- Lobster as a TL2 species; in terms of frequency, consumption is not culturally significant; TL4 specie is American eel – may be an ecological concern, but not an acute HH concern.
- Biogeochemical concerns due to wood waste – elevated % methylation on marshes; most concerning trophic transfer pathway to marsh species is via terrestrial food web for migratory songbirds.
- Surface sediment concentrations < 10 mg/kg in vicinity of site and < 1 mg/kg across majority of the estuary.
- No acute, severe or obvious human health concerns; species of greatest consumption are lobster and ducks, both of which have consumption restrictions in place via licensing structure/programs.

Wabigoon River, Ontario

- Glaciated terrain and very low sed. rates (~ 0.3 cm /yr)
- Walleye and Northern pike as TL4 species; consumption is culturally significant
- Mill effluent renders the river suboxic/anoxic in summer; stratification of an in-river lake contributes to > 2 ug/g in walleye.
- Surface sediment mercury concentrations exceed 50 mg/kg in vicinity of mill and are elevated consistently > 1 mg/kg for a distance of ~ 40 miles downstream.
- Human health impacts are acute, severe and with multi-generational manifestation; fish are consumed whether or not a consumption restriction is in place.

Grassy Narrows ANA Community – This is Living Downstream



Health in Grassy Narrows 'significantly worse' than other First Nations: report



Grassy Narrows ANA Community

Research

A Section 508-conformant HTML version of this article is available at <https://doi.org/10.1289/EHP11301>.

The Contribution across Three Generations of Mercury Exposure to Attempted Suicide among Children and Youth in Grassy Narrows First Nation, Canada: An Intergenerational Analysis

Donna Mergler,¹ Aline Philibert,¹ Myriam Fillion,^{2,1} and Judy Da Silva³



Mercury exposure and premature mortality in the Grassy Narrows First Nation community: a retrospective longitudinal study

Aline Philibert, Myriam Fillion, Donna Mergler

Summary

Background Little is known about the influence of toxic exposures on reduced life expectancy in First Nations people in Canada. The Grassy Narrows First Nation community have lived with the consequences of one of the worst environmental disasters in Canadian history. In the early 1960s, 10 000 kg of mercury (Hg) was released into their aquatic ecosystem. Although Hg concentration in fish, their dietary staple, decreased over time, it remains high. We aimed to examine whether elevated Hg exposure over time contributes to premature mortality (younger than 60 years) in this community.

Methods We did longitudinal and case-control analyses with data for individuals of the Grassy Narrows First Nation community. In 2019, the community obtained their historical Hg biomarker data from a government surveillance programme, which was then shared with the authors. A matched-pair approach allowed us to compare longitudinal hair Hg concentration between cases (individuals who died aged younger than 60 years) and controls (individuals who lived beyond 60 years). Matching criteria included year of birth (allowing 2 years either side), sex, and a minimum of four hair Hg concentration measures, of which at least two were in the same year. Analyses included change-point detection, interrupted time series, mixed models, and Cox survival models.

Findings We analysed data collected between Jan 1, 1970, and Jan 31, 1997, for 657 individuals (319 women and 338 men, born between 1884 and 1991) for whom we assembled a retrospective database of yearly measures of hair Hg concentration (n=3603). Hair Hg concentration decreased over time. A subgroup of 222 individuals (107 women and 115 men) reached or could have reached 60 years old by August, 2019. There was an increased risk of dying at a younger age among those with at least one hair Hg measure of 15 µg/g or more (adjusted hazard ratio 1.55, 95% CI 1.11–2.16; p=0.0088). Among the deceased individuals (n=154), longevity decreased by 1 year with every 6.25 µg/g (4.35–14.29) increase in hair Hg concentration. Analyses of 36 matched pairs showed that hair Hg concentration of those who died aged younger than 60 years was 4.7 µg/g higher (3.4–5.9) than controls.

Interpretation The consistent findings between our different analyses support an association between long-term Hg exposure from freshwater fish consumption and premature mortality in this First Nation community. There is a need to do risk-benefit analyses of freshwater fish consumption in environmentally contaminated regions.



Lancet Planet Health 2020; 4: e141-48

This online publication has been corrected. The corrected version first appeared at [thelancet.com/planetary-health](https://www.thelancet.com/planetary-health) on May 11, 2020, and further corrections have been made on July 15, 2020

Université du Québec à Montréal, Centre de recherche interdisciplinaire sur le bien-être, la santé, la société et l'environnement (Cinbiose), Montréal, QC, Canada (A Philibert PhD, Prof M Fillion PhD, Prof D Mergler PhD); and Département Science et Technologie, Université TÉLUQ, Montréal, QC, Canada (Prof M Fillion)

Correspondence to: Prof Donna Mergler, Centre de recherche interdisciplinaire sur le bien-être, la santé, la société et l'environnement (Cinbiose), Université du Québec à Montréal, Montréal, QC H3C 3P8, Canada mergler.donna@uqam.ca

CANADA

Ontario knew about Grassy Narrows mercury site for decades, but kept it secret

Toronto Star
November 11, 2017

A confidential 2016 report says provincial officials were told in the 1990s that the site of a paper mill near Grassy Narrows First Nation was contaminated with mercury — and that the poison is likely still present.

Walleye (1970 – 2017)




TABLE 7.1. Annual mercury discharges from the chlor-alkali plant, Great Lakes Forest Products Limited, Dryden.

Year	Mercury Loss (kg)
1962-69	1100
1970	350
1971	9.1
1972	2.3
1973	2.1
1974	1.7
1975	2.0
1976	1.2
1977	1.7





Figure 7.1 Mean monthly mercury concentrations of mill effluent, August, 197

Waste from mill worsening mercury contamination in river near Grassy Narrows: study



Industrial discharge from a paper mill in northern Ontario is exacerbating mercury contamination in a river system near a First Nation that has been plagued with mercury poisoning for decades, a new study suggests. Grassy Narrows Chief Rudy Turtle holds a sign as he marches with supporters through downtown Toronto in a 2019 handout photo.



Grassy Narrows chief calls out Ottawa for 'ridiculous' delays to mercury treatment centre construction

Trudeau said 'money is not the objection' to building the centre during 2019 election debate

 Brett Forester · CBC News · Posted: Feb 16, 2024 12:42 PM EST | Last Updated: February 16




Rudy Turtle, chief of Grassy Narrows First Nation, addresses a rally against mining proposals on First Nations territory in Toronto in July 2023. (Evan Mitsui/CBC)


Relationship between total mercury (Hg_T) and methyl mercury (MeHg) in **sediment**

- 1400 data pairs
- Salinity continuum
- Range in organic carbon concentration and quality
- Range in level of contamination
- Variable sources

Note that the relationship in this graph is descriptive of sampling conditions; it is not predictive of remedial response (meaning: we should be careful of trying to over-specify the extent to which decreasing Hg_T necessarily results in predictable declines in MeHg).



How Do These Data Distribute by Source/Type of Environment?



Note that the relationships in this graph are descriptive of sampling conditions; they are not predictive of remedial response (meaning: we should be careful of trying to overspecify the extent to which decreasing Hg_T necessarily results in predictable declines in MeHg).

THE LOOP CURRENT
 The Gulf's largest current, the Loop Current, enters from the Caribbean as the Yucatán Current. Running to depths of 2,600 feet, it can swing directly east to join the Gulf Stream or surge north before curling back through the Straits of Florida. If it penetrates deeply into the Gulf, it often sheds a great eddy, which drifts westward. The Loop Current could carry oil from a Gulf spill up the Atlantic coast.



- Map Legend**
- Coastal wetland
 - Coastal protected area
 - Active federal lease as of March 2010
 - Shallow water
 - Deep water (1,000 to 4,999 feet)
 - Ultradeep water (5,000 feet or more)
 - Oil or gas offshore platform
 - Oil or gas well
 - Crude oil or gas terminal
 - Oil refinery
 - Oil- or gas-related pipeline
 - Maritime boundary
 - Planning area boundary

NATIONAL GEOGRAPHIC

Gulf of Mexico

A GEOGRAPHY OF OFFSHORE OIL

For the past half century, oil has driven the economy of the Gulf of Mexico. A third of U.S. oil production flows from nearly 3,500 platforms in the Gulf, with thousands of miles of pipeline delivering oil and natural gas to shore. Since the first Gulf well was drilled off Louisiana in 1938, in less than 15 feet of water, close-in reserves have been depleted and exploration has marched off the continental shelf, onto the continental slope, and beyond. Today Gulf oil is deep oil; the bulk of U.S. production draws from wells in more than a thousand feet of water. U.S. Gulf oil reserves are estimated at 44.9 billion barrels, but as the *Deepwater Horizon* disaster showed, the challenges of deep drilling are formidable.

MEXICO'S OIL DROP
 Daily output of Pemex, the state-owned oil monopoly, hit 3.4 million barrels in 2004 but has fallen to 2.6 million. The drop is blamed on poor management and declining close-in reserves. Three-quarters of the oil comes from the Gulf, where Mexico has estimated reserves of at least 11.3 billion barrels. Oil and gas sales fund a third of the federal budget; the U.S. is the top importer.


IXTOC I OIL SPILL
 Mexico's Ixtoc I well blew out in the Bay of Campeche in 1979 and flowed for 295 days. Some 3.5 million barrels of oil fouled hundreds of miles of shore as far as Padre Island, Texas. Most habitat recovered, but three-inch-thick tar mats remain in some lagoons. Ixtoc I ranked as the world's largest accidental marine spill until it was surpassed by Macondo.

U.S. PLATFORMS
 A waterborne city of oil rigs rises off the coasts of Texas, Louisiana, Alabama, and Mississippi—but not Florida. Exploration off the Florida coast, starting in the 1940s, yielded tangle dry holes, and the potential threat to the state's top industry, tourism, has been considered too great a risk.

DEEPWATER DISASTER
 Working in 5,000 feet of water, the *Deepwater Horizon* rig had drilled 13,000 feet into the seabed when the exploratory Macondo well blew out on April 20, 2010. The explosion and fire sank the 58,000-ton mobile rig, killing 11 workers. An estimated 4.9 million barrels of oil flowed from the well, creating the worst accidental marine oil spill in history.




NEW DEPTHS
 The world's deepest offshore well, the Tiber well (art. above) reaches nearly six miles below the Gulf's seafloor. Not yet operational, it was drilled in

Important to remember that not all industrial facilities that may be sources of mercury to the environment are onshore.



CUBA'S POTENTIAL
 Estimates for Cuba's offshore oil reserves range as high as 4.6 billion barrels. No production has begun, but offshore development rights have been leased by the government to companies from Brazil, India, Malaysia, Norway, Spain, Venezuela, and Vietnam.

Mercury cell chlor-alkali process



Facility operations not uncommonly released
~ 10 tons of mercury into adjacent waters (plus
unquantified volumes into the atmosphere)

Locations of Former or Current Mercury Cell Chlor-Alkali Facilities

Acme, NC
 Ashtabula, OH
 Augusta, GA
 Bellingham Bay, WA
 Berlin, NH
 Brunswick, GA
 Calvert City, KY
 Charleston, TN
 Deer Park, TX
 Delaware City, DE
 East St. Louis, IL
 Geismar, LA
 Lake Charles, LA
 Lavaca Bay, TX
 Lemoyne, AL
 Linden, NJ
 McIntosh, AL
 Midland, MI
 Mobile, AL
 Moundsville, WV
 Muscle Shoals, AL
 New Castle, DE
 New Martinsville, WV
 Niagara Falls, NY (*2)
 Orrington, ME
 Plaquemine, LA
 Port Edwards, WI
 St. Gabriel, LA
 Syracuse, NY
 Dalhousie, NB
 Saguenay, Quebec
 Beauharnois, Quebec
 Marathon, ON
 Cornwall, ON
 Sarnia, ON
 Dryden, ON
 Port Abercrombie, NS
 Squamish, BC
 Coatzacoalcas-Minatitlán, Mexico
 García Nuevo León, Mexico
 Ecatepec de Morelos, Mexico
 Santa Clara, Mexico

Sagua la Grande, Cuba
 Managua, Nicaragua
 Cartagena, Colombia
 Maracaibo, Venezuela
 Lima, Peru
 Callao, Peru
 Itapagipe, Bahia, Brazil
 Botafogo River estuary, Brazil
 Ribeira Bay, Brazil
 Santo André, São Paulo, Brazil
 Santos- Cubatão, Brazil
 Acari-São João de Merití River, Brazil
 Cinco Saltos / Upper Negro River, Argentina
 Bahía Blanca, Argentina
 Argentina -3
 Montevideo, Uruguay

Bohus, Sweden
 Stenungsund, Sweden
 Skoghall, Sweden
 Domsjö, Sweden
 Koepmanholmen, Sweden
 Sweden-6
 Sarpsborg, Norway
 Kokemäenjoki, Finland
 Oulu, Finland
 Aetna, Finland
 Kuusankoski, Finland
 Runcorn, UK
 Sandbach, UK
 Staveley, UK
 Fleetwood, UK
 Ellesmere Port, UK
 Fermoy, Ireland
 Vieux-Thann, France
 Tavaux, France
 St. Auban, France
 Jarrie, France
 Loos, France
 Lavéra, France
 Mazingarbe, France
 Jemeppe -sur-Sambre, Belgium
 Antwerp, Belgium
 Tessenderlo, Belgium
 Botlek-Rotterdam, Netherlands
 Beek Geleen, Netherlands
 Delfzijl, Netherlands
 Linne Herten, Netherlands
 Hengelo, Netherlands
 Bitterfeld, GR
 Burghausen, GR
 Dormagen, GR
 Frankfurt, GR
 Gendorf, GR
 Gersthofen, GR
 Ibbenbüren, GR
 Hürth-Knapsack, GR
 Krefeld-Uerdingen, GR
 Lampertheim, GR
 Leverkusen, GR
 Ludwigshafen, GR
 Lüttsdorf, GR (*2)
 Marl, GR (*2)
 Markredwitz, GR
 Rheinfelden, GR
 Schkopau, GR
 Schkopau, GR
 Uerdingen, GR
 Wilhelmshafen, GR
 Hallein, Austria
 Brückl, Austria
 Neratovice, Czech Republic
 Ústí nad Labem, Czech Republic
 Pardubice, Czech Republic

Nováky, Slovakia
 Tamów, Poland
 Bydgoszcz, Poland
 Brzeg Dony, Poland
 Torrelavega, Spain
 Vilaseca, Spain
 Huelva, Spain
 Flix, Spain
 Jodar, Spain
 Monzon, Spain
 Hernani, Spain
 Sabinanigo/Huesca, Spain
 Pova de Santa Ir. , Portugal
 Ria de Aveiro, Portugal
 Pallanza Bay, Italy
 Priolo, Italy
 Augusta Bay, Italy
 Montova, Italy
 Tavazzano, Italy
 Gela, Italy
 Saline di Volterra Italy
 Rosignano Solvay, Italy
 Brescia, Italy
 Bussi, Italy
 Pieve Vergonte, Italy
 Volterra, Italy
 Toreviscosa, Italy
 Porto Marghera/Venice, Italy
 Ravenna, Italy
 Varna, Bulgaria
 Kazincbarcika, Hungary
 Ramnicu Valcea, Romania
 Vlora, Albania
 Thessaloniki, Greece
 Slovenia -1
 Serbia -1
 Aliaga, Turkey
 Kyiv, Ukraine
 Kalush, Ukraine (?)

Kirovo-Chepetsk, Russia
 Sterlitamak, Russia (possibly *2)
 Volgograd, Russia (possibly *2)
 Sayansk, Russia
 Ufa, Russia
 Dzerzhinsk, Russia
 Novodvinsk, Russia
 Chapaevsk, Russia
 Irkutsk, Russia
 Komsomolsk-on-Amur, Russia
 Usolye-Sibirskoye, Russia
 Koryazhma, Russia

Yavan, Tajikistan
 Sumgait, Azerbaijan
 Pavlodar, Kazakhstan
 Temirtau, Kazakhstan

Locations of Former or Current Mercury Cell Chlor-Alkali Facilities

Bandar Imam, Iran
 Kor River site, Iran (?)
 Iran -3
 Iraq -1
 Israel-1
 Syria-1

Alroli, India
 Mumbai, India
 Ganjam, India
 Singrauli, India
 Kota, Rajasthan, India
 Renukoot, Uttar Pradesh, India
 Nagda, Madhya Pradesh, India
 Kala Shah Kaku, Pakistan
 Myanmar -1
 Minamata Bay, Japan
 Niigata, Japan
 Omi, Japan
 Arai/Kosai, Japan
 Uto, Japan
 Shin-Nanyo, Yamaguchi, Japan
 Tokuyama, Yamaguchi, Japan
 Mizushima, Kurashiki, Okayama, Japan
 Takasago, Hyōgo, Japan
 Nobeoka City, Miyazaki, Japan
 Kashima, Ibaraki, Japan
 Ichihara City, Chiba, Japan
 An Ning, China
 Huludao, China
 Tianjin, China
 Yongjing, China
 Qingzhen, China
 Jilin City, China
 Dezhou City, Shandong Province, China
 Kaohsiung City, Taiwan
 Merak, Banten Province, Indonesia
 Indonesia-2
 Philippines-1
 Hamhung, North Korea
 North Korea-2

Alexandria, Egypt
 Mohammedia, Morocco
 Algeria-1
 Angola-1
 Libya - 1



Mercury Cell Chlor-Alkali Facility – identified
 Mercury Cell Chlor-Alkali Facility – newly added
 Mercury Chlor-Alkali Facility– not confirmed*
 Acetaldehyde Facility – identified

*Facility may have operated with an asbestos diaphragm

- The **Penobscot River** is the second largest river system in New England
- The estuary is:
 - ~20 miles long
 - 12 ft tidal range
- Seasonally variable discharge:
 - 5000 – 60,000 cfs
- Glaciated terrain and a long narrow river channel upgradient of Frankfort
- A mercury cell chlor-alkali facility operated in the estuary from 1967 – 2000
- Preceding history of wood products industry complicates remediation of the estuary and extends a recovery timeline to 70+ years








So, where is oceanic mercury coming from?






<input checked="" type="checkbox"/>	● Operating	6525
<input checked="" type="checkbox"/>	● Construction	421

● Permitted	254
● Pre-Permit	229
● Announced	181

Also...is this what a **tipping point** looks like...?



● Retired	2955
● Cancelled	3063
● Shelved	210
● Mothballed	166

And on a smaller (individually) but no less dangerous
(individually AND globally) scale....



(gold mining is extraordinarily dangerous for those who
have to feed their families this way....)




Serra Pelada, Brazil

Photographer:
Sebastiao Salgado
(1944 - 2025)

1986 – 1989 | gold mine



All sectors - Atmospheric emissions



Source	Amount (kg)
Artisanal and Small Scale Mining	837,658
Stationary Combustion of Coal	473,777
Nonferrous Metals Production	326,657
Cement Production	233,168
Waste from Products	146,938

A global ocean inventory of anthropogenic mercury based on water column measurements

Carl H. Lamborg¹, Chad R. Hammerschmidt², Katlin L. Bowman², Gretchen J. Swarr¹, Kathleen M. Munson¹, Daniel C. Ohnemus¹, Phoebe J. Lam¹, Lars-Eric Heimbürger³, Micha J. A. Rijkenberg⁴ & Mak A. Saito¹





Abstract Monomethylmercury (CH₃Hg) is the only form of mercury (Hg) known to biomagnify in food webs. Here we investigate factors driving methylated mercury [MeHg = CH₃Hg + (CH₃)₂Hg] production and degradation across the global ocean and uptake and trophic transfer at the base of marine food webs. We develop a new global 3-D simulation of MeHg in seawater and phyto/zooplankton within the Massachusetts Institute of Technology general circulation model. We find that high modeled MeHg concentrations in polar regions are driven by reduced demethylation due to lower solar radiation and colder temperatures. In the eastern tropical subsurface waters of the Atlantic and Pacific Oceans, the model results suggest that high MeHg concentrations are associated with enhanced microbial activity and atmospheric inputs of inorganic Hg. Global budget analysis indicates that upward advection/diffusion from subsurface ocean provides 17% of MeHg in the surface ocean. Modeled open ocean phytoplankton concentrations are relatively uniform because lowest modeled seawater MeHg concentrations occur in oligotrophic regions with the smallest size classes of phytoplankton, with relatively high uptake of MeHg and vice versa. Diatoms and synechococcus are the two most important phytoplankton categories for transferring MeHg from seawater to herbivorous zooplankton, contributing 35% and 25%, respectively. Modeled ratios of MeHg concentrations between herbivorous zooplankton and phytoplankton are 0.74–0.78 for picoplankton (i.e., no biomagnification) and 2.6–4.5 for eukaryotic phytoplankton. The spatial distribution of the trophic magnification factor is largely determined by the zooplankton concentrations. Changing ocean biogeochemistry resulting from climate change is expected to have a significant impact on marine MeHg formation and bioaccumulation.


Distribution of mercury species across a zonal section of the eastern tropical South Pacific Ocean (U.S. GEOTRACES GP16)


Katlin L. Bowman^a  , Chad R. Hammerschmidt^a , Carl H. Lamborg^{b 1} , Gretchen J. Swarr^b , Alison M. Agather^a 

Highlights

- Total mercury was enriched in the Peru upwelling region and up to 20% of the upwelling flux was as monomethyl-mercury.
- Subsurface maxima of monomethyl-mercury and dimethyl-mercury were found in oxic and suboxic water.
- Methylated mercury concentrations were greatest in the eastern part of the section underlying productive surface waters.
- Mercury was not elevated in a metal-rich hydrothermal vent plume extending 4000km west from the East Pacific Rise.
- Deep water below 2500m was enriched with Hg, especially in warm bottom waters in the eastern part of the section.







- Significant organic matter breakdown consumes dissolved oxygen (dO_2);
- Sulfate (SO_4^{2-}) + very low dO_2 increases activity of sulfate-reducing **bacteria** (SRB);
- SRB in the presence of inorganic mercury (Hg^{2+}) generate methyl mercury (CH_3Hg^+) as a **by-product of respiration**;
- CH_3Hg^+ is 100× more toxic than Hg^{2+} and is retained in biological tissue to a greater extent than Hg^{2+}

PHYSICAL

Thermohaline Circulation
(+ time since the Industrial
Revolution and significant
global increase in coal
combustion)


- ✦ = North Atlantic (+ Polar)
- ★ = Upwelling Zones and OMZ
- ★ = Photic Zone (CMZ)

BIOLOGICAL

Most significant
bioaccumulation step
is between water
column and uptake by
phytoplankton

CHEMICAL

Factors that contribute BOD
and result in O_2 consumption
(there's no shortage of SO_4^{2-});
consider this spatially



Think of this overview of
risk profiles as describing
regions of the global ocean
and considering residence
time (τ) in the oceans